

Comparison of lactulose, metronidazole and hepatic specific diet in controlling clinical signs in dogs with congenital extrahepatic portosystemic shunts: A randomized clinical trial

G. Serrano¹, N. Devriendt², H. de Rooster², D. Paepe²

¹Small Animal Department, Faculty of Veterinary Medicine, Ghent University, Merelbeke, Belgium; ²Small Animal Department, Faculty of Veterinary Medicine, Ghent University, Merelbeke, Belgium

Current medical management of dogs with congenital extrahepatic portosystemic shunt (cEHPSS) comprises the use of hepatic specific diet (HSD), lactulose and/or antibiotic therapy, aiming to control clinical signs of hepatic encephalopathy (HE). Meta-analysis of people suffering from type C HE disclosed that antibiotic and disaccharide therapy are equally effective in controlling HE symptoms.

A triple-arm randomized clinical trial was designed to compare the efficacy of different treatment combinations [HSD and lactulose (HSD + LACT), HSD and metronidazole (HSD + METRO), and solely HSD] in controlling the clinical signs in dogs suffering from cEHPSS. Dogs were not allowed to be receiving medical therapy at the time of inclusion. The allocated 'initial' treatment was continued for 4 weeks. Subsequently, all dogs received the combined treatment with HSD, LACT and METRO for an additional 2 weeks. Standardized questionnaires (assessing neurologic, gastrointestinal and urinary signs) were completed by the owners and a clinical score was calculated at diagnosis and at 4 and 6 weeks.

Thirty-six dogs were included, 12 were allocated to each group. Two dogs, 1 from HSD + LACT group and 1 from HSD + METRO groups were subsequently excluded due to concomitant diseases. Thirty-three dogs had their clinical scores available at the 4-weeks recheck and 27 dogs completed the trial until week 6 (10 HSD + LACT, 9 HSD + METRO, and 8 HSD). At diagnosis, the clinical scores were similar in all treatment groups ($P = 0.89$). Dogs in the "HSD + METRO" as well as the "HSD + LACT" group had a significant reduction in their clinical scores after the initial treatment compared to at diagnosis ($P = 0.03$ and $P = 0.002$, respectively). In contrast, no clinical improvement was observed in the HSD group ($P = 0.401$). At 6 weeks, a trend to improvement was noticed in the HSD group ($P = 0.06$), but no further improvement of the clinical score was detected in both other groups ($P = 1$).

Combination of HSD + LACT or HSD + METRO effectively reduces the clinical scores in cEHPSS dogs, while sole HSD therapy is not sufficient to control the same clinical signs.

Disclosures

No disclosures to report.